

Directions: Differentiate the functions.

1. $y = \sin(7x + \pi)$

$$\frac{dy}{dx} = \cos(7x + \pi) \cdot (7 + 0) = \boxed{7 \cos(7x + \pi)}$$

2. $z = \sqrt[3]{w^3 + 8} = (w^3 + 8)^{\frac{1}{3}}$

$$\frac{dz}{dw} = \frac{1}{3} (w^3 + 8)^{-\frac{2}{3}} (3w^2 + 0) = \frac{3w^2}{3(w^3 + 8)^{\frac{2}{3}}} = \boxed{\frac{w^2}{\sqrt[3]{(w^3 + 8)^2}}}$$

3. $y = \sec^2(x) = (\sec(x))^2$

$$\frac{dy}{dx} = 2(\sec(x))^{2-1} \sec(x) \tan(x) = \boxed{2 \sec^2(x) \tan(x)}$$

4. $y = \sec(x^2)$

$$\frac{dy}{dx} = \boxed{\sec(x^2) \tan(x^2) 2x}$$

5. $D_x [x e^{\tan(3x)+1}] = D_x [x] e^{\tan(3x)+1} + x D_x [e^{\tan(3x)+1}]$

$$= 1 \cdot e^{\tan(3x)+1} + x e^{\tan(3x)+1} D_x [\tan(3x) + 1]$$

$$= \boxed{e^{\tan(3x)+1} + x e^{\tan(3x)+1} (\sec^2(3x) \cdot 3 + 0)}$$

Directions: Differentiate the functions.

1. $y = \sqrt{5x+1} = (5x+1)^{\frac{1}{2}}$

$$\frac{dy}{dx} = \frac{1}{2} (5x+1)^{-\frac{1}{2}} (5+0) = \frac{5}{2(5x+1)^{\frac{1}{2}}} = \boxed{\frac{5}{2\sqrt{5x+1}}}$$

2. $y = \cos(x^2)$

$$\frac{dy}{dx} = -\sin(x^2) 2x = \boxed{-2x \sin(x^2)}$$

3. $y = \cos^2(x^2) = (\cos(x^2))^2$

$$\frac{dy}{dx} = 2(\cos(x^2))^{2-1} (-\sin(x^2) 2x) = \boxed{-4x \cos(x^2) \sin(x^2)}$$

4. $z = \tan\left(\frac{e^w}{w+1}\right)$

$$\frac{dz}{dw} = \sec^2\left(\frac{e^w}{w+1}\right) \frac{e^w(w+1) - e^w(1+0)}{(w+1)^2}$$

$$= \boxed{\sec^2\left(\frac{e^w}{w+1}\right) \frac{we^w}{(w+1)^2}}$$

5. $y = e^{\tan(3x)+x} + x^2$

$$\frac{dy}{dx} = D_x \left[e^{\tan(3x)+x} \right] + D_x \left[x^2 \right]$$

$$= e^{\tan(3x)+x} D_x \left[\tan(3x)+x \right] + 2x$$

$$= \boxed{e^{\tan(3x)+x} (\sec^2(3x) \cdot 3 + 1) + 2x}$$

Directions: Differentiate the functions.

1. $z = \sqrt{4w^2 + 16} = (4w^2 + 16)^{1/2}$

$$\frac{dz}{dw} = \frac{1}{2}(4w^2 + 16)^{-1/2} \cdot 8w = \frac{4w}{(4w^2 + 16)^{1/2}} = \frac{4w}{\sqrt{4w^2 + 16}}$$

2. $y = e^{x^2 - x}$

$$\frac{dy}{dx} = e^{x^2 - x} (2x - 1)$$

$$= \frac{2w}{\sqrt{w^2 + 4}}$$

3. $y = \sin(e^{x^2 - x})$

$$\frac{dy}{dx} = \cos(e^{x^2 - x}) D_x [e^{x^2 - x}] = \cos(e^{x^2 - x}) e^{x^2 - x} (2x - 1)$$

4. $y = (4x^5 \cos(x) + 1)^{10}$

$$\frac{dy}{dx} = 10(4x^5 \cos(x) + 1)^9 D_x [4x^5 \cos(x) + 1]$$

$$= 10(4x^5 \cos(x) + 1)^9 (20x^4 \cos(x) - 4x^5 \sin(x))$$

5. $D_x \left[\frac{e^{\tan(x)}}{x} \right] =$

$$= \frac{D_x [e^{\tan(x)}] \cdot x - e^{\tan(x)} D_x [x]}{x^2}$$

$$= \frac{e^{\tan(x)} \sec^2(x) \cdot x - e^{\tan(x)} \cdot 1}{x^2} = \frac{e^{\tan(x)} x \sec^2(x) - 1}{x^2}$$