


The Derivative of e^x

This chapter's goal is to find a derivative rule for the natural exponential function. We ask: If $f(x) = e^x$, what is $f'(x)$? We will answer this by working out the limit $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$.

Actually, we already know a little about this from Chapter 5. In Section 5.6 we found that the tangent to the graph of $f(x) = e^x$ at the point $(0, 1)$ has a slope of 1. (Fact 5.2 on page 93). This fact is illustrated on the right. It tells us that



$$\begin{aligned} 1 = f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{e^{0+h} - e^0}{h} = \lim_{h \rightarrow 0} \frac{e^h - 1}{h}. \end{aligned}$$

We will need this fact shortly. Note that it gives the value of a certain limit:

$$\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1. \tag{19.1}$$

Now let's find the derivative of $f(x) = e^x$ using the limit definition of $f'(x)$.


$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{e^{x+h} - e^x}{h} && \text{(definition of } f'(x)) \\ &= \lim_{h \rightarrow 0} \frac{e^x e^h - e^x}{h} && \text{(using } e^{x+h} = e^x e^h) \\ &= \lim_{h \rightarrow 0} \frac{e^x (e^h - 1)}{h} && \text{(factor out } e^x) \\ &= \lim_{h \rightarrow 0} e^x \cdot \lim_{h \rightarrow 0} \frac{e^h - 1}{h} && \text{(limit law)} \\ &= e^x \cdot 1 = e^x && \text{(Equation 19.1)} \end{aligned}$$

We have just found that if $f(x) = e^x$, then $f'(x) = e^x$. In other words, e^x is its own derivative! This is our latest derivative rule.

Rule 6 $D_x [e^x] = e^x$.

Geometrically, this new rule tells us that the tangent to the graph of $y = e^x$ at the point (x, e^x) has slope e^x . (See the diagram on the right.) The slope at the point (x, e^x) is as big as the point is high.

The fact that e^x is its own derivative is yet another indication of how special the natural exponential function e^x is, and why we place more importance on it than on other exponential functions a^x . The derivative of e^x is e^x . (As we will see, the derivative of, say, 2^x is **not** 2^x .)



You will often use this new rule in conjunction with other rules. For example, suppose we need to find the derivative of $x^5 - 3e^x + 1$. The answer comes from combining Rule 6 with rules 1–5:

$$\begin{aligned} D_x [x^5 - 3e^x + 1] &= D_x [x^5] - D_x [3e^x] + D_x [1] \\ &= 5x^4 - 3D_x [e^x] + 0 \\ &= 5x^4 - 3e^x \end{aligned}$$

Of course you will typically skip steps and get the answer immediately.

Be careful not to apply Rule 6 blindly. Notice that, for instance, $D_x [e^3] = 0$ because $e^3 \approx 2.71828^3 = 20.08555$ is a constant, and the derivative of a constant is zero. (Some students mistakenly write $D_x [e^3] = e^3$, or, even worse, $D_x [e^3] = 3e^2$. These are **wrong**. The first is a misapplication of Rule 6. The second is a misapplication of the power rule.)

Exercises for Chapter 19

Find the derivatives of the following functions in problems 1–6.

1. $f(x) = \sqrt{2}e^x + \sqrt{x}$
2. $f(x) = \frac{1}{x} - e^x + 3$
3. $w = z + e^2$
4. $y = e^{5+x}$ Hint: $e^{a+b} = e^a e^b$.
5. $f(x) = 6x^3 + e^x - 4$
6. $f(x) = \frac{3}{x^4} + \frac{e^x}{3}$
7. Find the equation of the tangent line to $y = 3e^x$ at the point $(2, 3e^2)$.
8. For what x is the tangent to $y = e^x - x$ at $(x, e^x - x)$ horizontal?

Exercise Solutions for Chapter 19

$$\begin{aligned}
 \mathbf{1.} \quad D_x[\sqrt{2}e^x + \sqrt{x}] &= D_x[\sqrt{2}e^x] + D_x[\sqrt{x}] = \sqrt{2}D_x[e^x] + D_x[x^{1/2}] \\
 &= \sqrt{2}e^x + \frac{1}{2}x^{1/2-1} = \sqrt{2}e^x + \frac{1}{2}x^{-1/2} = \sqrt{2}e^x + \frac{1}{2x^{1/2}} = \boxed{\sqrt{2}e^x + \frac{1}{2\sqrt{x}}}
 \end{aligned}$$

$$\mathbf{3.} \quad \frac{d}{dz}[z + e^2] = 1 + 0 = \boxed{1}$$

$$\mathbf{5.} \quad f'(x) = 18x^2 + e^x$$

7. Find the equation of the tangent line to $y = 3e^x$ at the point $(2, 3e^2)$.

The slope of the tangent to $y = 3e^x$ at $(x, 3e^x)$ is $\frac{dy}{dx} = 3e^x$. We are interested in the tangent line at $(2, 3e^2)$, and its slope is $\left.\frac{dy}{dx}\right|_{x=2} = 3e^2$. So its slope is $m = 3e^2$ and it passes through $(2, 3e^2)$. We can get its equation with the point-slope formula.

$$\begin{aligned}
 y - y_0 &= m(x - x_0) \\
 y - 3e^2 &= 3e^2(x - 2) \\
 y &= 3e^2x - 3e^2 \cdot 2 + 3e^2 \\
 y &= 3e^2x - 3e^2
 \end{aligned}$$

Answer: The tangent line has equation $y = 3e^2x - 3e^2$.